

Europäisches Patentamt  
European Patent Office  
Office européen des brevets



(11) **EP 1 209 233 A2**

(12) **EUROPEAN PATENT APPLICATION**

(43) Date of publication:  
29.05.2002 Bulletin 2002/22

(51) Int Cl.7: **C12N 15/57, C12N 9/54,  
C12N 1/21, C11D 3/386**

(21) Application number: **01127851.2**

(22) Date of filing: **22.11.2001**

(84) Designated Contracting States:  
**AT BE CH CY DE DK ES FI FR GB GR IE IT LI LU  
MC NL PT SE TR**  
Designated Extension States:  
**AL LT LV MK RO SI**

(30) Priority: **22.11.2000 JP 2000355166  
12.04.2001 JP 2001114048**

(71) Applicant: **Kao Corporation  
Tokyo 103-8210 (JP)**

(72) Inventors:  
• **Hatada, Yuji**  
**Mihara-shi, Hiroshima 723-0052 (JP)**  
• **Ogawa, Akinori**  
**Haga-gun, Tochigi 321-3497 (JP)**

- **Kageyama, Yasushi**  
**Haga-gun, Tochigi 321-3497 (JP)**
- **Sato, Tsuyoshi**  
**Haga-gun, Tochigi 321-3497 (JP)**
- **Araki, Hiroyuki**  
**Haga-gun, Tochigi 321-3497 (JP)**
- **Sumitomo, Nobuyuki**  
**Haga-gun, Tochigi 321-3497 (JP)**
- **Okuda, Mitsuyoshi**  
**Haga-gun, Tochigi 321-3497 (JP)**
- **Saeki, Katsuhisa**  
**Haga-gun, Tochigi 321-3497 (JP)**

(74) Representative: **VOSSIUS & PARTNER  
Siebertstrasse 4  
81675 München (DE)**

(54) **Alkaline proteases**

(57) Provided are an alkaline protease wherein an amino acid residue at (a) position 84, (b) position 104, (c) position 256 or (d) position 369 of SEQ ID NO:1 or at a position corresponding thereto has been deleted or selected from: at position (a): an arginine residue, at position (b): a proline residue, at position (c): an alanine, serine, glutamine, valine, leucine, asparagine, glutamic acid or aspartic acid residue, and at position (d): an aspartic acid residue; an alkaline protease wherein an amino acid residue at (e) position 66 or 264, (f) position 57, each of 101 to 106, 136, 193 or 342, (g) position 46 or 205, (h) position 54, 119, 138, 148 or 195, (i) position 247, (j) position 124, (k) position 107 or (l) position 257 has been deleted or selected from: at position (e): a glutamine, aspartic acid or like residue, at position (f): a lysine, serine or like residue, at position (g): a tyrosine

or tryptophan residue, at position (h): a tryptophan, phenylalanine or like residue, at position (i): a tryptophan, phenylalanine or like residue, at position (j): an alanine or lysine residue, at position (k): a lysine, arginine or like residue, and at position (l): a valine or isoleucine residue; a gene encoding the alkaline protease; a recombinant vector containing the gene, a transformant containing the recombinant vector; and a detergent composition containing the alkaline protease.

The present invention makes it possible to provide alkaline proteases having activity even in the presence of a high concentration of fatty acids, having high specific activity and detergency and being useful as an enzyme to be incorporated in a detergent.

EP 1 209 233 A2

# Description

[0001] The present invention relates to alkaline proteases having high specific activity and strong oxidant resistance and as an enzyme to be added to a detergent, having excellent detergency.

[0002] Proteases have conventionally been used in a variety of fields such as various detergents including laundry detergents, cosmetics, bath agents, food modifiers, and pharmaceuticals such as digestion aids and anti-inflammatory agents. Among them, proteases for detergents are industrially produced in the largest amount and have a great market scale. A number of proteases are now put on the market.

[0003] In most cases, stains on clothes contain not only proteins but also plural components such as lipids and solid particles. There is accordingly a demand for detergents having detergency high enough to remove such actual complex stains. Finding, from such a viewpoint, alkaline proteases capable of retaining caseinolytic activity even in the presence of a high concentration of fatty acids and exhibiting excellent detergency even if the stain is composed of not a simple protein component but plural components, for example, protein and lipid, and having a molecular weight of about 43,000, the present inventors applied a patent (WO99/18218) on them.

[0004] Alkaline proteases superior to the above-described ones in specific activity, oxidant resistance and detergency and usable for detergents of wide-ranging compositions have however been requested.

[0005] The present inventors searched for such alkaline proteases mainly from enzyme variants. The above-described alkaline proteases are however utterly different in enzymological properties from serine proteases typified by subtilisin so that the modified site of subtilisin did not provide them with useful information. As a result of a further investigation, the present inventors have found that in order to obtain novel alkaline proteases having improved specific activity, stability against an oxidant and detergency while maintaining the properties of the above-described alkaline proteases, they must have a specific amino acid residue at a predetermined position of their amino acid sequence.

[0006] In one aspect of the present invention, there is thus provided an alkaline protease wherein an amino acid residue at (a) position 84, (b) position 104, (c) position 256 or (d) position 369 of SEQ ID NO:1 or at a position corresponding thereto has been deleted or selected from:

at position (a): an arginine residue,  
at position (b): a proline residue,  
at position (c): an alanine, serine, glutamine, valine, leucine, asparagine, glutamic acid or aspartic acid residue, and  
at position (d): an aspartic acid residue.

[0007] In another aspect of the present invention, there is also provided an alkaline protease wherein an amino acid residue at (e) position 66 or 264, (f) position 57, each of 101 to 106, 136, 193 or 342, (g) position 46 or 205, (h) position 54, 119, 138, 148 or 195, (i) position 247, (j) position 124, (k) position 107 or (l) position 257 of SEQ ID NO:1, or at a position corresponding thereto has been deleted or selected from:

at position (e): a glutamine, aspartic acid, serine, glutamic acid, alanine, threonine, leucine, methionine, cysteine, valine, glycine or isoleucine residue  
at position (f): a lysine, serine, glutamine, phenylalanine, valine, arginine, tyrosine, leucine, isoleucine, threonine, methionine, cysteine, tryptophan, aspartic acid, glutamic acid, histidine, proline or alanine residue,  
at position (g): a tyrosine, tryptophan, alanine, asparagine, glutamic acid, threonine, valine, leucine, isoleucine, histidine, serine, lysine, glutamine, methionine or cysteine residue,  
at position (h): a tryptophan, phenylalanine, alanine, asparagine, glutamic acid, threonine, valine, histidine, serine, lysine, glutamine, methionine, glycine, aspartic acid, proline, arginine or cysteine residue,  
at position (i): a tryptophan, phenylalanine, alanine, asparagine, glutamic acid, threonine, valine, leucine, isoleucine, histidine, serine, glutamine, methionine or cysteine residue,  
at position (j): an alanine or lysine residue,  
at position (k): a lysine, arginine, alanine or serine residue, and  
at position (l): a valine or isoleucine residue.

[0008] In a further aspect of the present invention, there are also provided a gene encoding the alkaline protease, a recombinant vector containing the gene and a transformant containing the vector.

[0009] In a still further aspect of the present invention, there is also provided a detergent composition containing the alkaline protease.

FIG. 1 is a diagram illustrating detergency of an alkaline protease variant;  
FIG. 2 is a diagram illustrating relative specific activity of each of alkaline protease variants; and FIG. 3 is a diagram illustrating relative residual activity of each of alkaline protease variants after treatment with an oxidant.

[0010] As described above, in the alkaline proteases of the present invention, an amino acid residue at (a) position 84, (b) position 104, (c) position 256 or (d) position 369 of SEQ ID NO:1 or at a position corresponding thereto has been deleted or selected from: at position (a): an arginine residue, at position (b): a proline residue, at position (c): an alanine, serine, glutamine, valine, leucine, asparagine, glutamic acid or aspartic acid residue, and at position (d): an aspartic acid residue; or an amino acid residue at (e) position 66 or 264, (f) position 57, each of 101 to 106, 136, 193 or 342, (g) position 46 or 205, (h) position 54, 119, 138, 148 or 195, (i) position 247, (j) position 124, (k) position 107 or (l) position 257 of SEQ ID NO:1 or at a position corresponding thereto has been deleted or selected from: at position (e): a glutamine, aspartic acid, serine, glutamic acid, alanine, threonine, leucine, methionine, cysteine, valine, glycine or isoleucine residue, at position (f): a lysine, serine, glutamine, phenylalanine, valine, arginine, tyrosine, leucine, isoleucine, threonine, methionine, cysteine, tryptophan, aspartic acid, glutamic acid, histidine, proline or alanine residue, at position (g): a tyrosine, tryptophan, alanine, asparagine, glutamic acid, threonine, valine, leucine, isoleucine, histidine, serine, lysine, glutamine, methionine or cysteine residue, at position (h): a tryptophan, phenylalanine, alanine, asparagine, glutamic acid, threonine, valine, histidine, serine, lysine, glutamine, methionine, glycine, aspartic acid, proline, arginine or cysteine residue, at position (i): a tryptophan, phenylalanine, alanine, asparagine, glutamic acid, threonine, valine, leucine, isoleucine, histidine, serine, glutamine, methionine or cysteine residue, at position (j): an alanine or lysine residue, at position (k): a lysine, arginine, alanine or serine residue, and at position (l): a valine or isoleucine residue.

[0011] Specifically, the alkaline proteases according to the present invention mean alkaline proteases having an amino acid sequence represented by SEQ ID NO:1 wherein the amino acid residue at a position selected from the above-described (a) to (d) and (e) to (l) has been deleted or predetermined, or another alkaline protease wherein the amino acid residue at a position corresponding thereto has been deleted or predetermined. They may be wild type enzymes, wild type variants or artificial variants.

[0012] The "another alkaline protease" may be a wild type enzyme or a wild type variant. That having oxidant resistance and having a molecular weight, as determined by SDS-PAGE, of  $43,000 \pm 2,000$  is preferred, of which that having an amino acid sequence showing at least 60% homology with the amino acid sequence of SEQ ID NO:1 is more preferred. Particularly preferred is that having an amino acid sequence showing at least 60% homology with the amino acid sequence of SEQ ID NO:1, having oxidant resistance, works on the alkaline side (pH 8 or greater), is stable with at least 80% residual activity when treated at pH 10 for 10 minutes even at 50°C, is inhibited by diisopropyl fluorophosphate (DFP) and phenylmethane sulfonyl fluoride (PMSF) and has a molecular weight, as determined by SDS-PAGE, of  $43,000 \pm 2,000$ . The term "having oxidant resistance" as used herein means that it has at least 50% of residual activity (synthetic substrate assay) when treated in a 50 mM hydrogen peroxide solution (containing 5 mM calcium chloride) at pH 10 (a 20 mM Britton-Robinson buffer) at 20°C for 20 minutes.

[0013] Examples of the "alkaline protease having an amino acid sequence represented by SEQ ID NO:1" include KP43 [derived from *Bacillus* sp. strain KSM-KP43 (FERM BP-6532), WO99/18218], while those of the "alkaline protease having an amino acid sequence showing at least 60% homology with the amino acid sequence of SEQ ID NO:1" include protease KP9860 having an amino acid sequence represented by SEQ ID NO:2 [derived from *Bacillus* sp. strain KSM-KP9860 (FERM BP-6534), WO99/18218], Protease E-1 having an amino acid sequence represented by SEQ ID NO:3 [derived from *Bacillus* sp. strain No. D-6 (FERM P-1592), JP740710], Protease Ya having an amino acid sequence represented by SEQ ID NO:4 [derived from *Bacillus* sp. strain Y (FERM BP-1029), JP861210], Protease SD521 having an amino acid sequence represented by SEQ ID NO:5 [derived from *Bacillus* sp. strain SD-521 (FERM BP-11162), JP910821], Protease A-1 having an amino acid sequence represented by SEQ ID NO:6 (derived from NCIB12289, WO8801293), and Protease A-2 having an amino acid sequence represented by SEQ ID NO:7 (derived from NCIB12513, WO8801293). Of these, the amino acid sequences selected from SEQ ID NOS. 2 to 7 or alkaline proteases showing at least 80%, more preferably at least 90%, especially at least 95% homology therewith are preferred.

[0014] The homology of an amino acid sequence is calculated by Lipman-Pearson's method (Science, **227**, 1435 (1985)).

[0015] The "amino acid residue at a corresponding position" can be identified by comparing amino acid sequences by using known algorithm, for example, that of Lipman-Pearson. The position of the "amino acid residue at a corresponding position" in the sequence of each protease can be determined by aligning the amino acid sequence of the protease in such a manner. It is presumed that the corresponding position exists at the three-dimensionally same position in the amino acid sequence of SEQ ID NO:1 and the amino acid residue existing at the same position brings about similar effects for a specific function of the protease.

[0016] Described specifically,

(a) the amino acid residue at position 84 of SEQ ID NO:1 is a lysine residue. By employing the above-described method, the amino acid residue at the position corresponding thereto can be identified as the lysine residue at position 83 of SEQ ID NO:3. This amino acid residue is preferably arginine.

(b) Although the amino acid residue at position 104 of SEQ ID NO:1 is a leucine residue, this amino acid residue

or an amino acid residue corresponding thereto is preferably a proline residue.

(c) Although the amino acid residue at position 256 of SEQ ID NO:1 is a methionine residue, particularly preferred as this amino acid residue is an alanine, serine, glutamine, valine, leucine, asparagine, glutamic acid or aspartic acid residue.

(d) Although the amino acid residue at position 369 of SEQ ID NO:1 is an aspartic acid residue, this amino acid residue or amino acid residue corresponding thereto is preferably an asparagine residue.

(e) Although the amino acid residue at position 66 or 264 of SEQ ID NO:1 is an asparagine residue, this amino acid residue is preferably a glutamine, aspartic acid, serine, glutamic acid, alanine, threonine, leucine, methionine, cysteine, valine, glycine or isoleucine residue, with an aspartic acid, serine or glutamic acid residue being particularly preferred. More preferred is the case wherein the amino acid residue at position 66 is an aspartic acid residue and that at position 264 is a serine residue.

(f) Although the amino acid residue at each of positions 57, 101 to 106, 136, 193 and 342 of SEQ ID NO:1 is a glycine residue, this amino acid residue is preferably a lysine, serine, glutamine, phenylalanine, valine, arginine, tyrosine, leucine, isoleucine, threonine, methionine, cysteine, tryptophan, aspartic acid, glutamic acid, histidine, proline or alanine residue. Particularly preferred is the case wherein the amino acid residue at position 57, 136, 193 or 342 is an alanine residue, or that at position 103 is an arginine residue.

(g) Although the amino acid residue at position 46 or 205 of SEQ ID NO:1 is a phenylalanine residue, this amino acid residue is preferably a tyrosine, tryptophan, alanine, asparagine, glutamic acid, threonine, valine, leucine, isoleucine, histidine, serine, lysine, glutamine, methionine or cysteine residue. Particularly preferred is the case wherein the amino acid residue at position 46 is a leucine residue.

(h) Although the amino acid residue at position 54, 119, 138, 148 or 195 of SEQ ID NO:1 is a tyrosine residue, this amino acid residue is preferably a tryptophan, phenylalanine, alanine, asparagine, glutamic acid, threonine, valine, histidine, serine, glutamine, methionine, glycine, aspartic acid, proline, lysine, arginine or cysteine residue. Particularly preferred is the case wherein the amino acid residue at position 195 is an alanine, aspartic acid, glutamic acid, glutamine, valine, tryptophan, glycine, lysine, threonine, methionine, cysteine, phenylalanine, proline, serine, arginine, asparagine or histidine residue.

(i) Although the amino acid residue at position 247 of SEQ ID NO:1 is a lysine residue, this amino acid residue is preferably a tryptophan, phenylalanine, alanine, asparagine, glutamic acid, threonine, valine, leucine, isoleucine, histidine, serine, glutamine, methionine or cysteine residue. As the amino acid residue at position 247, an arginine or threonine residue is particularly preferred.

(j) Although the amino acid residue at position 124 of SEQ ID NO:1 is an arginine residue, this amino acid residue is preferably an alanine or lysine residue.

(k) Although the amino acid residue at position 107 of SEQ ID NO:1 is a leucine residue, this amino acid residue is preferably a lysine, arginine, alanine or serine residue, with a lysine residue being particularly preferred.

(l) Although the amino acid residue at position 257 of SEQ ID NO:1 is an alanine residue, this amino acid residue is preferably a valine or isoleucine residue, with a valine residue being particularly preferred.

[0017] With regards to "another alkali protease" which is preferred among the above-exemplified ones, positions corresponding to (a) to (d) and (e) to (l) of the amino acid sequence (SEQ ID NO:1) of Protease KP43 and specific examples of an amino acid residue are shown below (Table 1-a, Table 1-b).

Table 1-a

Position	Proteases						
	TS43 SEQ ID NO:1	9860 SEQ ID NO:2	E-1 SEQ ID NO:3	Ya SEQ ID NO:4	SD-521 SEQ ID NO:5	A-1 SEQ ID NO:6	A-2 SEQ ID NO:7
(a)	84Lys	84Lys	83Lys	83Lys	83Lys	84Lys	83Lys
(b)	104Leu	104Leu	103Leu	103Leu	103Leu	104Leu	103Leu
(c)	256Met	256Met	255Met	255Met	255Met	256Met	255Met
(d)	369Asp	369Asp	368Asp	368Asp	368Asp	369Asp	368Asp

Table 1-b

Position	Proteases						
	TS43 SEQ ID NO:1	9860 SEQ ID NO:2	E-1 SEQ ID NO:3	Ya SEQ ID NO:4	SD-521 SEQ ID NO:5	A-1 SEQ ID NO:6	A-2 SEQ ID NO:7
(e)	66Asn 264Asn	66Asn 264Asn	66Asn 263Asn	66Asn 263Asn	66Asn 263Asn	66Asn 264Asn	66Asn 263Asn
(f)	57Gly 101Gly 102Gly 103Gly 105Gly 106Gly 136Gly 193Gly 342Gly	57Gly 101Ser 102Gly 103Gly 105Gly 106Gly 136Gly 193Gly 342Gly	56Gly 100Ser 101Gly 102Gly 104Gly 105Gly 135Gly 192Gly 341Gly	56Gly 100Ser 101Gly 102Gly 104Gly 105Gly 135Gly 192Gly 341Gly	56Gly 100Ser 101Gly 102Gly 104Gly 105Gly 135Gly 192Gly 341Gly	57Gly 101Asn 102Gly 103Gly 105Gly 106Gly 136Gly 193Gly 342Gly	56Gly 100Gly 101Gly 102Gly 104Gly 105Gly 135Gly 192Gly 341Gly
(g)	46Phe 205Phe	46Phe 205Phe	46Phe 204Phe	46Phe 204Phe	46Phe 204Phe	46Phe 205Phe	46Phe 204Phe
(h)	195Tyr	195Tyr	194Ile	194Ile	194Leu	195Tyr	194Tyr
(i)	247Lys	247Lys	246Lys	246Lys	246Lys	247Lys	246Lys
(j)	124Arg	124Arg	123Arg	123Arg	123Arg	124Arg	123Arg
(k)	107Leu	107Leu	106Leu	106Leu	106Leu	107Leu	106Leu
(l)	257Ala	257Ala	256Ala	256Ala	256Ala	257Ala	256Ala

[0018] In the alkaline proteases of the present invention, deletion of an amino acid residue or selection in (a) to (d) or (e) to (l) may be conducted at two or more positions simultaneously.

[0019] When the alkaline protease of the present invention is a variant, the "protease having an amino acid sequence represented by SEQ ID NO:1" or the above-exemplified "another alkaline protease" serves as an alkaline protease prior to mutation (which may be called "parent alkaline protease"). By introducing mutation to a desired site of this parent alkaline protease, the alkaline protease of the present invention is available. For example, it is available by deleting or substituting, with another amino acid residue, the amino acid residue at a position selected from the above-described (a) to (d) and (e) to (l) of the amino acid sequence of SEQ ID NO:1 of Protease KP43 or at the corresponding position of the amino acid sequence of another alkaline protease, more specifically, amino acid sequence represented by SEQ ID NOS:2 to 7.

[0020] The alkaline protease of the present invention can be obtained, for example, by introducing mutation to a cloned gene encoding a parent alkaline protease, transforming a proper host by using the resulting mutated gene and then culturing the recombinant host. Cloning of the gene for encoding a parent alkaline protease may be carried out using an ordinary gene recombination technique, for example, in accordance with the process as described in WO99/18218, JP901128 or WO98/56927.

[0021] For mutation of a gene encoding a parent alkaline protease, either one of random mutation or site-specific mutation which is prevalent now can be adopted. More specifically, mutation can be effected using, for example, "Site-Directed Mutagenesis System Mutan-Super Express Kit" of Takara Shuzo Co., Ltd. By using recombinant PCR (polymerase chain reaction) as described in "PCR protocols" (Academic Press, New York, 1990), a desired sequence of a gene can be replaced with a sequence of another gene corresponding to the desired sequence.

[0022] For production of the protease variant of the present invention by using the resulting mutated gene, usable is, for example, the following process. A DNA encoding the protease variant of the present invention is stably amplified by linking the mutated gene with a DNA vector capable of amplifying it stably or by introducing the mutated gene onto a chromosomal DNA capable of maintaining it stably and then, the gene is introduced into a host permitting stable and

efficient expression of the gene, whereby the variant protease is produced. Hosts satisfying the above-described conditions include microorganisms belonging to *Bacillus* sp., *Escherichia coli*, mold, yeast and *Actinomyces*.

[0023] The alkaline protease of the present invention thus obtained has stable protease activity in an alkaline region, is free from the inhibition of caseinolytic activity by higher fatty acids, and has a molecular weight, as determined by SDS-PAGE,  $43,000 \pm 2,000$ . For example, the protease variant available from, as a parent strain, the protease having an amino acid sequence of SEQ ID NO:1 has the below-described physicochemical properties.

(i) Acting pH range

[0024] It acts in a wide pH range of from 4 to 13 and exhibits at least 80% of the optimum pH active value at pH 6 to 12.

(ii) Stable pH range

[0025] It is stable within a pH range of 6 to 11 under the treating conditions at 40°C for 30 minutes.

(iii) Influence of fatty acids

[0026] Its caseinolytic activity is not inhibited by oleic acid.

[0027] Such proteases of the present invention have excellent specific activity, oxidant resistance and detergency and are therefore useful as an enzyme to be incorporated in various detergent compositions. Particularly, the proteases wherein the amino acid residue at position (a) to (d) of SEQ ID NO:1 or at a position corresponding thereto has been deleted or specified are superior in detergency. Among them, those having, as the amino acid residue at (c) position 256 or at a position corresponding thereto, an alanine, serine, glutamine, valine, leucine, asparagine, glutamic acid or aspartic acid residue have both high specific activity and strong oxidant resistance. The proteases wherein the amino acid residue at position (e) to (l) of SEQ ID NO:1 or at a position corresponding thereto has been deleted or specified have particularly excellent specific activity.

[0028] The above-described protease may be added to the detergent composition of the present invention in an amount sufficient to permit exhibition of its activity. Although 0.1 to 5000 P.U. can be added per 1 kg of the detergent composition, 1000 P.U. or less, preferably 500 P.U. is added in consideration of economy.

[0029] To the detergent composition of the present invention, various enzymes can be used in combination with the alkaline protease of the present invention. Examples include hydrolases, oxidases, reductases, transferases, lyases, isomerases, ligases and synthetases. Of these, proteases, cellulases, lipases, keratinases, esterases, cutinases, amylases, pullulanases, pectinases, mannanases, glucosidases, glucanases, cholesterol oxidases, peroxidases, laccases and proteases other than the alkaline protease used in the present invention are preferred.

[0030] Proteases include commercially available Alcalase, Esperase, Savinase and Everlase (each, product of Novo Nordisk), Properase and Purafect (each, product of Genencor International Inc.), and KAP (Kao Corp). Cellulases include Cellzyme and Carezyme (each, product of Novo Nordisk), KAC (Kao Corp.) and alkaline cellulase produced by *Bacillus* sp. strain KSM-S237 as described in Japanese Patent Application Laid-Open No. Hei 10-313859. Amylases include Termamyl and Duramyl (each, product of Novo Nordisk), Purastar (Genencor International Inc.), and KAM (Kao Corp.). Lipases include Lipolase and Liporase Ultra (each, product of Novo Nordisk). The above-exemplified enzyme may be incorporated in an amount of 0.001 to 10%, preferably 0.03 to 5%.

[0031] A surfactant may be incorporated in an amount of 0.5 to 60 wt.% (which will hereinafter be called "%", simply) in the detergent composition. To a powdery detergent composition and a liquid detergent composition, addition of 10 to 45% and 20 to 50% are preferred, respectively. When the detergent composition of the present invention is a bleaching detergent or automatic dishwasher detergent, the surfactant may usually be added in an amount of 1 to 10%, preferably 1 to 5%.

[0032] A divalent metal ion scavenger may be added in an amount of 0.01 to 50%, with 5 to 40% being preferred.

[0033] An alkali agent and inorganic salt may be added in an amount of 0.01 to 80%, preferably 1 to 40%.

[0034] An antisoil redeposition agent may be added in an amount of 0.001 to 10%, preferably 1 to 5%.

[0035] A bleaching agent (ex. hydrogen peroxide or percarbonate) is added preferably in an amount of 1 to 10%. Upon use of the bleaching agent, 0.01 to 10% of a bleaching activator can be added.

[0036] As a fluorescent brightener, biphenyl type ones (such as "Tinopal CBS-X") and stilbene type ones (such as DM fluorescent dye) can be used. It is added preferably in an amount of 0.001 to 2%.

[0037] The detergent composition of the present invention can be prepared in a conventional manner by using the alkaline protease obtained by the above-described process and the above-described known detergent components in combination. The detergent form can be selected according to the using purpose. Examples include liquid, powder and granule.

[0038] When the alkaline protease of the present invention is added to a powdery detergent composition, it is pre-

ferred to prepare detergent particles in advance and then mix therein the alkaline protease granules in accordance with the process as described in Japanese Patent Application Laid-Open No. Sho 62-25790 in order to avoid the contact of workers or end users with the enzyme upon preparation or use of the detergent or to prevent heat-induced deactivation or decomposition of the enzyme.

**[0039]** The detergent composition of the present invention thus available is usable as a laundry detergent, bleaching detergent, automatic dishwasher detergent, pipe cleaner and artificial tooth cleaner. Use as a laundry detergent, bleaching detergent or automatic dishwasher detergent is particularly preferred.

#### Example 1

**[0040]** Mutation was introduced at random into a protease structural gene of about 2.0 kb including a termination codon by the following manner. First, PCR was conducted using a primer capable of amplifying this 2.0kb. A PCR master mix contained 5 ng of a template DNA, 20 pmol of a phosphorylated primer, 20 nmol of each dNTP, 1  $\mu$ mol of Tris/HCl (pH 8.3), 5  $\mu$ mol of KCl, 0.15  $\mu$ mol of  $MgCl_2$  and 2.5U TaqDNA polymerase, and its total amount was adjusted to 100  $\mu$ L. After modification of the template by allowing it to stand at 94°C for 5 minutes, PCR was performed for 30 cycles, each cycle consisting of treatment at 94°C for 1 min, at 55°C for 1 min and at 72°C for 1.5 min. The PCR product was purified by "PCR product purification Kit" (product of Boeringer Mannheim), followed by elution in 100  $\mu$ L of sterile water. With 1  $\mu$ L of the eluate, second PCR was conducted under conditions similar to those of the first PCR except for the template DNA. After completion of the second PCR, the PCR product was purified in a similar manner to the first PCR, followed by elution in 100  $\mu$ L of sterile water.

**[0041]** The amplified DNA fragment was integrated in a vector by polymerase reaction using "LATAq" produced by Takara Shuzo Co., Ltd. Described specifically, after addition of 5  $\mu$ L of a buffer for LATAq (a 10-fold concentrate), 8  $\mu$ L of a dNTP solution and 0.5  $\mu$ L of LATAq DNA polymerase, and as a template, 20 ng of plasmid pHA64TS (having a protease structural gene linked with an expression vector pHA64) to 35  $\mu$ L of the purified eluate, the total amount was adjusted to 50  $\mu$ L. PCR reaction of the resulting liquid was carried out for 30 cycles, each consisting of treatment at 94°C for 1 min, 55°C for 1 min and 72°C for 4 min. By the subsequent ethanol precipitation, the PCR product was collected. This PCR product had a shape of a plasmid having a nick at the 5' prime end of the primer. Ligase reaction by T4 ligase (product of Takara Shuzo Co., Ltd.) was conducted to link this nick portion.

**[0042]** By using 10  $\mu$ L of this ligase reaction mixture, transformation of the *Bacillus subtilis* strain ISW1214 was conducted, whereby about  $4 \times 10^5$  transformants were obtained. The resulting transformants of the strain ISW1214 were cultured on a skin-milk-containing medium (containing 1% skim milk, 1% bactotrypton, 1% sodium chloride, 0.5% yeast extract, 1.5% agar and 7.5  $\mu$ g/ml of tetracycline) and halo formation, which was presumed to reflect the protease secretion amount, was observed.

#### Example 2: Purification of an enzyme

**[0043]** The protease active fraction was prepared in the following manner. The transformants obtained in Example 1 was cultured at 30°C for 60 hours on a medium A (3% polypeptone S (product of Nippon Pharmaceutical), 0.5% yeast extract, 1% fish meat extract (product of Wako Pure Chemical Industries, Ltd.), 0.15% dipotassium phosphate, 0.02% magnesium sulfate 7 hydrate, 4% maltose and 7.5  $\mu$ g/mL of tetracycline). The supernatant of the thus-obtained cultured medium was added with ammonium sulfate to give 90% saturation, whereby salting-out of protein was caused. The sample obtained by salting-out was dissolved in a 10 mM tris HCl buffer (pH 7.5) containing 2 mM of calcium chloride. The resulting solution was dialyzed overnight against the same buffer by using a dialysis membrane. The fraction in the dialysis membrane was applied to DEAE Bio-Gel A (product of Bio-Rad Laboratories) equilibrated with a 10 mM tris HCl buffer (pH 7.5) containing 2 mM calcium chloride to collect the protease active fraction not adsorbed to the ion-exchanger. This active fraction was applied further to "SP-Toyopearl 550W" (product of Tosoh Corp.) equilibrated with the same buffer, followed by elution with a 0 to 50 mM sodium chloride solution, whereby a protease active fraction was obtained. The resulting fraction was analyzed by SDS-PAGE electrophoresis to confirm that the protease was obtained as substantially uniform protein. The protein concentration was measured in accordance with the method of Lowry, et al. (J. Biol. Chem. **193**, 265-275(1981)) by using bovine serum albumin (product of Bio-Rad Laboratories) as a standard.

#### Example 3: Measuring method of protease activity

##### (1) Synthetic substrate assay

**[0044]** A decomposition rate was measured using a synthetic peptide made of Glt-Ala-Ala-Pro-Leu(A-A-P-L) as a substrate. Described specifically, a 50 mM borate/KCl buffer (pH 10.5) containing each enzyme to be evaluated and

## EP 1 209 233 A2

3 mM of Glt-A-A-P-L-pNA (product of Peptide Institute, Inc) was kept at 30°C for 10 minutes and then, an absorbance at 420 nm was periodically measured. The peptide hydrolyzing activity was determined from an increasing ratio of the absorbance at 420 nm per unit hour. The protein was determined using a protein assay kit of Bio-Rad Laboratories.

### (2) Natural substrate assay

[0045] After 1.0 mL of a 50 mM borate buffer (pH 10) containing 1% (w/v) of casein was kept at 30°C for 5 minutes, 0.1 mL of an enzyme solution was added and reaction was conducted for 15 minutes. To the reaction mixture, 2.0 mL of a reaction-stopping solution (0.11M trichloroacetic acid - 0.22M sodium acetate - 0.33M acetic acid) was added. The resulting mixture was allowed to stand at room temperature for 30 minutes and the filtered. The acid soluble protein in the filtrate was quantitatively determined by the modified method of Lowry, et al. Described specifically, after addition of 2.5 mL of an alkaline copper solution [1% potassium sodium tartrate : 1% copper sulfate : 1% sodium carbonate = 1:1:100] to the filtrate, the resulting solution was allowed to stand at room temperature for 10 minutes. Then, 0.25 mL of a diluted phenol solution (a phenol reagent (product of Kanto Kagaku) diluted 2-fold with ion exchange water) was added. After the resulting mixture was kept at 30°C for 30 minutes, absorbance at 660 nm was measured. One enzyme unit was designated as a quantity of the enzyme for liberating the acid soluble protein hydrolysis product corresponding to 1 mmol of tyrosine for 1 min in the above-described reaction.

### Example 4

#### (1) Preparation of granular detergent

[0046] Detergency of the detergent as described in Example 3 of WO99/29830 was evaluated. Described specifically, 465 kg of water was poured in a mixing tank of 1 m<sup>3</sup> equipped with a stirring blade. After its water temperature reached 55°C, 48 kg of a 50% (w/v) aqueous solution of sodium dodecylbenzenesulfonate and 135 kg of a 40% (w/v) aqueous solution of sodium polyacrylate were added. After stirring for 15 minutes, 120 kg of sodium carbonate, 60 kg of sodium sulfate, 9 kg of sodium sulfite and 3 kg of a fluorescent dye were added. After stirring for further 15 minutes, 300 kg of zeolite was added. The mixture was stirred for 30 minutes to yield a uniform slurry (the slurry had a water content of 50 wt.%). By spraying this slurry from a pressure spraying nozzle disposed in the vicinity of the top of a spray drying tower, base granules were obtained (a high temperature gas was fed to the spray drying tower at 225°C from the tower bottom and discharged from the tower top at 105°C).

[0047] Then, 15 parts by weight of a nonionic surfactant, 15 parts by weight of a 50 wt.% aqueous solution of sodium dodecylbenzenesulfonate and 1 part by weight of polyethylene glycol were mixed under heating to 70°C, whereby a mixture was obtained. In a Loedige mixer (product of Matsuzaka Giken Co., Ltd., capacity: 20L, equipped with a jacket), 100 parts by weight of the base granules obtained above were charged and stirring by a main shaft (150 rpm) and chopper (4000 rpm) was started. Warm water of 75°C was caused to flow in the jacket at 10 L/min, the mixture was charged therein in 3 minutes, and then stirring was conducted for 5 minutes. The surface of the detergent particles were covered with 10 parts by weight of crystalline aluminosilicate, whereby the final product of the granular detergent was obtained.

#### [Raw materials used]

Aqueous solution of sodium dodecylbenzenesulfonate: "Neopelex F65" (product of Kao Corp.)

[0048] Nonionic surfactant: "Emulgen 108KM" (product of Kao) added with 8.5 moles, on average, of ethylene oxide  
Aqueous solution of sodium polyacrylate: having an average molecular weight of 10000 (prepared in accordance with the process as described in Example of Japanese Patent Publication No. Hei 2-24283)

Sodium carbonate: dense ash (product of Central Glass Co., Ltd.)

Zeolite: "Zeolite 4A" having an average particle size of 3.5 µm (product of Tosoh Corp)

Polyethylene glycol: "K-PEG6000" (average molecular weight of 8500, product of Kao Corp.)

Fluorescent dye: "Tinopai CBS-X" (product of Ciba Geigy)

#### (2) Preparation of granulated protease

[0049] From the alkaline proteases of the present invention and a purified preparation of a parent alkaline protease, granulated protease was prepared based on the process as described in Japanese Patent Application Laid-Open No. Sho 62-257990 (6P.U./g)

(3) Measurement of detergency

[0050] In 1L of an aqueous calcium solution (71.2 mg calcium carbonate/1L) adjusted to 20°C, 0.67 g of each of the detergent compositions as shown in Table 2 was dissolved. With the resulting solution, a test cloth ("EMPA117" - (prepared by Swiss Federal Laboratories for Materials Testing and Research, blood/milk/carbon) cut into a piece of 6 × 6 cm was washed using a Terg-O-tometer (product of Ueshima Seisakusho) at 20°C and 100 rpm for 10 minutes. After rinsing and drying, the brightness was measured using a spectrophotometer ("CM3500d", product of MINOLTA). A detergency was calculated based on the below-described equation. The results are shown in Table 2.

Detergency (%) =  $\frac{\text{Brightness of the test fabric after washing} - \text{that before washing}}{\text{Brightness of the test fabric before soiling} - \text{that before washing}} \times 100$

[0051] Measuring results of the detergency of the protease variants obtained in Example 1 are shown in FIG. 1. The alkaline protease variants of the present invention each exhibited superior detergency to wild type enzymes to which mutation had not been introduced.

Table 2

			Invention products					Comparative products	
			1	2	3	4	5	1	2
Parts by weight	Granulated alkaline proteases of this invention	K84R	0.5	0.5	0.5	0.5	0.5		
		L104P							
		M256S							
M256A									
D369N									
	Granulated parent alkaline protease							0.5	
	Granular detergent	99.5						100	
Detergency (%)			38	38	36	36	34	31	23

Example 5

[0052] Measuring results, in accordance with the synthetic substrate assay or natural substrate assay, of protease activity of the protease variants obtained in Example 1 (the proteases modified at 195-position and 256-position amino acid residues, respectively was measured by the latter assay, while the other proteases were measured by the former assay) are shown in FIG. 2. The alkaline protease variants of the present invention exhibited high specific activity.

Example 6

[0053] In 2 mL of a 100 mM borate buffer (pH 10.5) containing 3% of aqueous hydrogen peroxide, a 50 µl portion of each of the protease variants obtained by purification in Example 1 was added. The resulting mixture was allowed to stand at 30°C for 30 minutes. After addition of an adequate amount of catalase (product of Boehringer Mannheim) to remove excess hydrogen peroxide, the residual protease activity was measured by the synthetic substrate assay. In FIG. 3, the residual activity after treatment with aqueous hydrogen peroxide is shown relative to the activity before treatment set at 100%.

[0054] The alkaline protease variants of the present invention exhibited higher oxidant resistance than the parent alkaline protease.

[0055] The present invention makes it possible to provide alkaline proteases having activity even under a high concentration of fatty acids, having high specific activity, oxidant resistance and detergency, and being useful as an enzyme to be incorporated in a detergent.

# EP 1 209 233 A2

## SEQUENCE LISTING

<110> KAO CORPORATION

5 <120> Alkaline protease

<130> F 2935 EP

<150> JP P2000-355166

10 <151> 2000-11-22

<150> JP P2001-114048

<151> 2001-4-12

15 <160> 7

<170> PatentIn Ver. 2.1

<210> 1

20 <211> 434

<212> PRT

<213> Bacillus sp.

<400> 1

25 Asn Asp Val Ala Arg Gly Ile Val Lys Ala Asp Val Ala Gln Ser Ser  
1 5 10 15

Tyr Gly Leu Tyr Gly Gln Gly Gln Ile Val Ala Val Ala Asp Thr Gly  
20 25 30

30 Leu Asp Thr Gly Arg Asn Asp Ser Ser Met His Glu Ala Phe Arg Gly  
35 40 45

Lys Ile Thr Ala Leu Tyr Ala Leu Gly Arg Thr Asn Asn Ala Asn Asp  
50 55 60

35 Thr Asn Gly His Gly Thr His Val Ala Gly Ser Val Leu Gly Asn Gly  
65 70 75 80

40 Ser Thr Asn Lys Gly Met Ala Pro Gln Ala Asn Leu Val Phe Gln Ser  
85 90 95

Ile Met Asp Ser Gly Gly Gly Leu Gly Gly Leu Pro Ser Asn Leu Gln  
100 105 110

45 Thr Leu Phe Ser Gln Ala Tyr Ser Ala Gly Ala Arg Ile His Thr Asn  
115 120 125

Ser Trp Gly Ala Ala Val Asn Gly Ala Tyr Thr Thr Asp Ser Arg Asn  
130 135 140

50 Val Asp Asp Tyr Val Arg Lys Asn Asp Met Thr Ile Leu Phe Ala Ala  
145 150 155 160

Gly Asn Glu Gly Pro Asn Gly Gly Thr Ile Ser Ala Pro Gly Thr Ala  
165 170 175

55

EP 1 209 233 A2

5 Lys Asn Ala Ile Thr Val Gly Ala Thr Glu Asn Leu Arg Pro Ser Phe  
 180 185 190  
 Gly Ser Tyr Ala Asp Asn Ile Asn His Val Ala Gln Phe Ser Ser Arg  
 195 200 205  
 10 Gly Pro Thr Lys Asp Gly Arg Ile Lys Pro Asp Val Met Ala Pro Gly  
 210 215 220  
 Thr Phe Ile Leu Ser Ala Arg Ser Ser Leu Ala Pro Asp Ser Ser Phe  
 225 230 235 240  
 15 Trp Ala Asn His Asp Ser Lys Tyr Ala Tyr Met Gly Gly Thr Ser Met  
 245 250 255  
 Ala Thr Pro Ile Val Ala Gly Asn Val Ala Gln Leu Arg Glu His Phe  
 260 265 270  
 20 Val Lys Asn Arg Gly Ile Thr Pro Lys Pro Ser Leu Leu Lys Ala Ala  
 275 280 285  
 Leu Ile Ala Gly Ala Ala Asp Ile Gly Leu Gly Tyr Pro Asn Gly Asn  
 290 295 300  
 25 Gln Gly Trp Gly Arg Val Thr Leu Asp Lys Ser Leu Asn Val Ala Tyr  
 305 310 315 320  
 30 Val Asn Glu Ser Ser Ser Leu Ser Thr Ser Gln Lys Ala Thr Tyr Ser  
 325 330 335  
 Phe Thr Ala Thr Ala Gly Lys Pro Leu Lys Ile Ser Leu Val Trp Ser  
 340 345 350  
 35 Asp Ala Pro Ala Ser Thr Thr Ala Ser Val Thr Leu Val Asn Asp Leu  
 355 360 365  
 Asp Leu Val Ile Thr Ala Pro Asn Gly Thr Gln Tyr Val Gly Asn Asp  
 370 375 380  
 40 Phe Thr Ser Pro Tyr Asn Asp Asn Trp Asp Gly Arg Asn Asn Val Glu  
 385 390 395 400  
 Asn Val Phe Ile Asn Ala Pro Gln Ser Gly Thr Tyr Thr Ile Glu Val  
 405 410 415  
 45 Gln Ala Tyr Asn Val Pro Val Gly Pro Gln Thr Phe Ser Leu Ala Ile  
 420 425 430  
 50 Val Asn  
 <210> 2  
 <211> 434  
 <212> PRT  
 55 <213> Bacillus sp.

EP 1 209 233 A2

<400> 2

5	Asn	Asp	Val	Ala	Arg	Gly	Ile	Val	Lys	Ala	Asp	Val	Ala	Gln	Ser	Ser	1	5	10	15
	Tyr	Gly	Leu	Tyr	Gly	Gln	Gly	Gln	Ile	Val	Ala	Val	Ala	Asp	Thr	Gly	20	25	30	
10	Leu	Asp	Thr	Gly	Arg	Asn	Asp	Ser	Ser	Met	His	Glu	Ala	Phe	Arg	Gly	35	40	45	
	Lys	Ile	Thr	Ala	Leu	Tyr	Ala	Leu	Gly	Arg	Thr	Asn	Asn	Ala	Asn	Asp	50	55	60	
15	Thr	Asn	Gly	His	Gly	Thr	His	Val	Ala	Gly	Ser	Val	Leu	Gly	Asn	Gly	65	70	75	80
	Ala	Thr	Asn	Lys	Gly	Met	Ala	Pro	Gln	Ala	Asn	Leu	Val	Phe	Gln	Ser	85	90	95	
20	Ile	Met	Asp	Ser	Ser	Gly	Gly	Leu	Gly	Gly	Leu	Pro	Ser	Asn	Leu	Gln	100	105	110	
	Thr	Leu	Phe	Ser	Gln	Ala	Phe	Ser	Ala	Gly	Ala	Arg	Ile	His	Thr	Asn	115	120	125	
25	Ser	Trp	Gly	Ala	Ala	Val	Asn	Gly	Ala	Tyr	Thr	Thr	Asp	Ser	Arg	Asn	130	135	140	
30	Val	Asp	Asp	Tyr	Val	Arg	Lys	Asn	Asp	Met	Thr	Ile	Leu	Phe	Ala	Ala	145	150	155	160
	Gly	Asn	Glu	Gly	Pro	Asn	Gly	Gly	Thr	Ile	Ser	Ala	Pro	Gly	Thr	Ala	165	170	175	
35	Lys	Asn	Ala	Ile	Thr	Val	Gly	Ala	Thr	Glu	Asn	Leu	Arg	Pro	Ser	Phe	180	185	190	
	Gly	Ser	Tyr	Ala	Asp	Asn	Ile	Asn	His	Val	Ala	Gln	Phe	Ser	Ser	Arg	195	200	205	
40	Gly	Pro	Thr	Lys	Asp	Gly	Arg	Ile	Lys	Pro	Asp	Val	Met	Ala	Pro	Gly	210	215	220	
	Thr	Tyr	Ile	Leu	Ser	Ala	Arg	Ser	Ser	Leu	Ala	Pro	Asp	Ser	Ser	Phe	225	230	235	240
45	Trp	Ala	Asn	His	Asp	Ser	Lys	Tyr	Ala	Tyr	Met	Gly	Gly	Thr	Ser	Met	245	250	255	
50	Ala	Thr	Pro	Ile	Val	Ala	Gly	Asn	Val	Ala	Gln	Leu	Arg	Glu	His	Phe	260	265	270	
	Val	Lys	Asn	Arg	Gly	Ile	Thr	Pro	Lys	Pro	Ser	Leu	Leu	Lys	Ala	Ala	275	280	285	

55

EP 1 209 233 A2

5 Leu Ile Ala Gly Ala Ala Asp Val Gly Leu Gly Tyr Pro Asn Gly Asn  
 290 295 300  
 Gln Gly Trp Gly Arg Val Thr Leu Asp Lys Ser Leu Asn Val Ala Tyr  
 305 310 315 320  
 10 Val Asn Glu Ser Ser Ala Leu Ser Thr Ser Gln Lys Ala Thr Tyr Thr  
 325 330 335  
 Phe Thr Ala Thr Ala Gly Lys Pro Leu Lys Ile Ser Leu Val Trp Ser  
 340 345 350  
 15 Asp Ala Pro Ala Ser Thr Thr Ala Ser Val Thr Leu Val Asn Asp Leu  
 355 360 365  
 Asp Leu Val Ile Thr Ala Pro Asn Gly Thr Arg Tyr Val Gly Asn Asp  
 370 375 380  
 20 Phe Ser Ala Pro Phe Asp Asn Asn Trp Asp Gly Arg Asn Asn Val Glu  
 385 390 395 400  
 Asn Val Phe Ile Asn Ser Pro Gln Ser Gly Thr Tyr Thr Ile Glu Val  
 405 410 415  
 25 Gln Ala Tyr Asn Val Pro Val Gly Pro Gln Asn Phe Ser Leu Ala Ile  
 420 425 430  
 30 Val Asn  
 <210> 3  
 <211> 433  
 <212> PRT  
 <213> Bacillus sp.  
 35 <400> 3  
 Asn Asp Val Ala Arg Gly Ile Val Lys Ala Asp Val Ala Gln Asn Asn  
 1 5 10 15  
 40 Tyr Gly Leu Tyr Gly Gln Gly Gln Val Val Ala Val Ala Asp Thr Gly  
 20 25 30  
 Leu Asp Thr Gly Arg Asn Asp Ser Ser Met His Glu Ala Phe Arg Gly  
 35 40 45  
 45 Lys Ile Thr Ala Leu Tyr Ala Leu Gly Arg Thr Asn Asn Ala Asn Asp  
 50 55 60  
 Pro Asn Gly His Gly Thr His Val Ala Gly Ser Val Leu Gly Asn Ala  
 65 70 75 80  
 50 Leu Asn Lys Gly Met Ala Pro Gln Ala Asn Leu Val Phe Gln Ser Ile  
 85 90 95  
 55 Met Asp Ser Ser Gly Gly Leu Gly Gly Leu Pro Ser Asn Leu Asn Thr

EP 1 209 233 A2

	100	105	110
5	Leu Phe Ser Gln Ala Trp Asn Ala Gly Ala Arg Ile His Thr Asn Ser 115 120 125		
	Trp Gly Ala Pro Val Asn Gly Ala Tyr Thr Ala Asn Ser Arg Gln Val 130 135 140		
10	Asp Glu Tyr Val Arg Asn Asn Asp Met Thr Val Leu Phe Ala Ala Gly 145 150 155 160		
	Asn Glu Gly Pro Asn Ser Gly Thr Ile Ser Ala Pro Gly Thr Ala Lys 165 170 175		
15	Asn Ala Ile Thr Val Gly Ala Thr Glu Asn Tyr Arg Pro Ser Phe Gly 180 185 190		
20	Ser Ile Ala Asp Asn Pro Asn His Ile Ala Gln Phe Ser Ser Arg Gly 195 200 205		
	Ala Thr Arg Asp Gly Arg Ile Lys Pro Asp Val Thr Ala Pro Gly Thr 210 215 220		
25	Phe Ile Leu Ser Ala Arg Ser Ser Leu Ala Pro Asp Ser Ser Phe Trp 225 230 235 240		
	Ala Asn Tyr Asn Ser Lys Tyr Ala Tyr Met Gly Gly Thr Ser Met Ala 245 250 255		
30	Thr Pro Ile Val Ala Gly Asn Val Ala Gln Leu Arg Glu His Phe Ile 260 265 270		
35	Lys Asn Arg Gly Ile Thr Pro Lys Pro Ser Leu Ile Lys Ala Ala Leu 275 280 285		
	Ile Ala Gly Ala Thr Asp Val Gly Leu Gly Tyr Pro Ser Gly Asp Gln 290 295 300		
40	Gly Trp Gly Arg Val Thr Leu Asp Lys Ser Leu Asn Val Ala Tyr Val 305 310 315 320		
	Asn Glu Ala Thr Ala Leu Thr Thr Gly Gln Lys Ala Thr Tyr Ser Phe 325 330 335		
45	Gln Thr Gln Ala Gly Lys Pro Leu Lys Ile Ser Leu Val Trp Thr Asp 340 345 350		
	Ala Pro Gly Ser Thr Thr Ala Ser Tyr Thr Leu Val Asn Asp Leu Asp 355 360 365		
50	Leu Val Ile Thr Ala Pro Asn Gly Gln Lys Tyr Val Gly Asn Asp Phe 370 375 380		
55	Ser Tyr Pro Tyr Asp Asn Asn Trp Asp Gly Arg Asn Asn Val Glu Asn 385 390 395 400		

EP 1 209 233 A2

5 Val Phe Ile Asn Ala Pro Gln Ser Gly Thr Tyr Thr Ile Glu Val Gln  
405 410 415

Ala Tyr Asn Val Pro Ser Gly Pro Gln Arg Phe Ser Leu Ala Ile Val  
420 425 430

10 His

<210> 4  
<211> 433  
<212> PRT  
<213> Bacillus sp.

15 <400> 4  
Asn Asp Val Ala Arg Gly Ile Val Lys Ala Asp Val Ala Gln Asn Asn  
1 5 10 15

20 Tyr Gly Leu Tyr Gly Gln Gly Gln Val Val Ala Val Ala Asp Thr Gly  
20 25 30

Leu Asp Thr Gly Arg Asn Asp Ser Ser Met His Glu Ala Phe Arg Gly  
35 40 45

25 Lys Ile Thr Ala Leu Tyr Ala Leu Gly Arg Thr Asn Asn Ala Ser Asp  
50 55 60

Pro Asn Gly His Gly Thr His Val Ala Gly Ser Val Leu Gly Asn Ala  
65 70 75 80

30 Leu Asn Lys Gly Met Ala Pro Gln Ala Asn Leu Val Phe Gln Ser Ile  
85 90 95

35 Met Asp Ser Ser Gly Gly Leu Gly Gly Leu Pro Ser Asn Leu Asn Thr  
100 105 110

Leu Phe Ser Gln Ala Trp Asn Ala Gly Ala Arg Ile His Thr Asn Ser  
115 120 125

40 Trp Gly Ala Pro Val Asn Gly Ala Tyr Thr Ala Asn Ser Arg Gln Val  
130 135 140

Asp Glu Tyr Val Arg Asn Asn Asp Met Thr Val Leu Phe Ala Ala Gly  
145 150 155 160

45 Asn Glu Gly Pro Asn Ser Gly Thr Ile Ser Ala Pro Gly Thr Ala Lys  
165 170 175

Asn Ala Ile Thr Val Gly Ala Thr Glu Asn Tyr Arg Pro Ser Phe Gly  
180 185 190

50 Ser Ile Ala Asp Asn Pro Asn His Ile Ala Gln Phe Ser Ser Arg Gly  
195 200 205

55 Ala Thr Arg Asp Gly Arg Ile Lys Pro Asp Val Thr Ala Pro Gly Thr

EP 1 209 233 A2

	210	215	220
5	Phe Ile Leu Ser Ala Arg Ser Ser Leu Ala Pro Asp Ser Ser Phe Trp 225 230 235 240		
	Ala Asn Tyr Asn Ser Lys Tyr Ala Tyr Met Gly Gly Thr Ser Met Ala 245 250 255		
10	Thr Pro Ile Val Ala Gly Asn Val Ala Gln Leu Arg Glu His Phe Ile 260 265 270		
	Lys Asn Arg Gly Ile Thr Pro Lys Pro Ser Leu Ile Lys Ala Ala Leu 275 280 285		
15	Ile Ala Gly Ala Thr Asp Val Gly Leu Gly Tyr Pro Asn Gly Asp Gln 290 295 300		
20	Gly Trp Gly Arg Val Thr Leu Asn Lys Ser Leu Asn Val Ala Tyr Val 305 310 315 320		
	Asn Glu Ala Thr Ala Leu Ala Thr Gly Gln Lys Ala Thr Tyr Ser Phe 325 330 335		
25	Gln Ala Gln Ala Gly Lys Pro Leu Lys Ile Ser Leu Val Trp Thr Asp 340 345 350		
	Ala Pro Gly Ser Thr Thr Ala Ser Tyr Thr Leu Val Asn Asp Leu Asp 355 360 365		
30	Leu Val Ile Thr Ala Pro Asn Gly Gln Lys Tyr Val Gly Asn Asp Phe 370 375 380		
35	Ser Tyr Pro Tyr Asp Asn Asn Trp Asp Gly Arg Asn Asn Val Glu Asn 385 390 395 400		
	Val Phe Ile Asn Ala Pro Gln Ser Gly Thr Tyr Ile Ile Glu Val Gln 405 410 415		
40	Ala Tyr Asn Val Pro Ser Gly Pro Gln Arg Phe Ser Leu Ala Ile Val 420 425 430		
	His		
45	<210> 5 <211> 433 <212> PRT <213> Bacillus sp.		
50	<400> 5 Asn Asp Val Ala Arg Gly Ile Val Lys Ala Asp Val Ala Gln Asn Asn 1 5 10 15		
	Tyr Gly Leu Tyr Gly Gln Gly Gln Val Val Ala Val Ala Asp Thr Gly 20 25 30		
55			

EP 1 209 233 A2

	Leu	Asp	Thr	Gly	Arg	Asn	Asp	Ser	Ser	Met	His	Glu	Ala	Phe	Arg	Gly	
			35					40					45				
5	Lys	Ile	Thr	Ala	Leu	Tyr	Ala	Leu	Gly	Arg	Thr	Asn	Asn	Ala	Asn	Asp	
		50					55					60					
	Pro	Asn	Gly	His	Gly	Thr	His	Val	Ala	Gly	Ser	Val	Leu	Gly	Asn	Ala	
		65				70					75					80	
10	Leu	Asn	Lys	Gly	Met	Ala	Pro	Gln	Ala	Asn	Leu	Val	Phe	Gln	Ser	Ile	
					85					90					95		
	Met	Asp	Ser	Ser	Gly	Gly	Leu	Gly	Gly	Leu	Pro	Ser	Asn	Leu	Asn	Thr	
15				100					105					110			
	Leu	Phe	Ser	Gln	Ala	Trp	Asn	Ala	Gly	Ala	Arg	Ile	His	Thr	Asn	Ser	
			115					120					125				
20	Trp	Gly	Ala	Pro	Val	Asn	Gly	Ala	Tyr	Thr	Ala	Asn	Ser	Arg	Gln	Val	
		130					135					140					
	Asp	Glu	Tyr	Val	Arg	Asn	Asn	Asp	Met	Thr	Val	Leu	Phe	Ala	Ala	Gly	
		145				150					155					160	
25	Asn	Glu	Gly	Pro	Asn	Ser	Gly	Thr	Ile	Ser	Ala	Pro	Gly	Thr	Ala	Lys	
					165					170					175		
	Asn	Ala	Ile	Thr	Val	Gly	Ala	Thr	Glu	Asn	Tyr	Arg	Pro	Ser	Phe	Gly	
				180					185					190			
30	Ser	Leu	Ala	Asp	Asn	Pro	Asn	His	Ile	Ala	Gln	Phe	Ser	Ser	Arg	Gly	
			195					200					205				
	Ala	Thr	Arg	Asp	Gly	Arg	Ile	Lys	Pro	Asp	Val	Thr	Ala	Pro	Gly	Thr	
35			210				215					220					
	Phe	Ile	Leu	Ser	Ala	Arg	Ser	Ser	Leu	Ala	Pro	Asp	Ser	Ser	Phe	Trp	
		225				230				235						240	
40	Ala	Asn	Tyr	Asn	Ser	Lys	Tyr	Ala	Tyr	Met	Gly	Gly	Thr	Ser	Met	Ala	
					245					250					255		
	Thr	Pro	Ile	Val	Ala	Gly	Asn	Val	Ala	Gln	Leu	Arg	Glu	His	Phe	Ile	
				260					265					270			
45	Lys	Asn	Arg	Gly	Ile	Thr	Pro	Lys	Pro	Ser	Leu	Ile	Lys	Ala	Ala	Leu	
			275					280					285				
	Ile	Ala	Gly	Ala	Thr	Asp	Val	Gly	Leu	Gly	Tyr	Pro	Ser	Gly	Asp	Gln	
		290				295						300					
50	Gly	Trp	Gly	Arg	Val	Thr	Leu	Asp	Lys	Ser	Leu	Asn	Val	Ala	Tyr	Val	
		305				310					315					320	
55	Asn	Glu	Ala	Thr	Ala	Leu	Ala	Thr	Gly	Gln	Lys	Ala	Thr	Tyr	Ser	Phe	

EP 1 209 233 A2

				325					330					335		
5	Gln	Ala	Gln	Ala	Gly	Lys	Pro	Leu	Lys	Ile	Ser	Leu	Val	Trp	Thr	Asp
				340					345					350		
	Ala	Pro	Gly	Ser	Thr	Thr	Ala	Ser	Tyr	Thr	Leu	Val	Asn	Asp	Leu	Asp
			355					360					365			
10	Leu	Val	Ile	Thr	Ala	Pro	Asn	Gly	Gln	Lys	Tyr	Val	Gly	Asn	Asp	Phe
		370					375					380				
	Ser	Tyr	Pro	Tyr	Asp	Asn	Asn	Trp	Asp	Gly	Arg	Asn	Asn	Val	Glu	Asn
15		385				390				395					400	
	Val	Phe	Ile	Asn	Ala	Pro	Gln	Ser	Gly	Thr	Tyr	Thr	Ile	Glu	Val	Gln
				405					410					415		
	Ala	Tyr	Asn	Val	Pro	Ser	Gly	Pro	Gln	Arg	Phe	Ser	Leu	Ala	Ile	Val
20				420					425					430		
	His															
	<210>	6														
25	<211>	434														
	<212>	PRT														
	<213>	Bacillus sp.														
	<400>	6														
30	Asn	Asp	Val	Ala	Arg	Gly	Ile	Val	Lys	Ala	Asp	Val	Ala	Gln	Ser	Ser
	1				5					10					15	
	Tyr	Gly	Leu	Tyr	Gly	Gln	Gly	Gln	Val	Val	Ala	Val	Ala	Asp	Thr	Gly
			20						25					30		
35	Leu	Asp	Thr	Gly	Arg	Asn	Asp	Ser	Ser	Met	His	Glu	Ala	Phe	Arg	Gly
		35						40					45			
	Lys	Ile	Thr	Ala	Ile	Tyr	Ala	Leu	Gly	Arg	Thr	Asn	Asn	Ala	Asn	Asp
40		50					55					60				
	Pro	Asn	Gly	His	Gly	Thr	His	Val	Ala	Gly	Ser	Val	Leu	Gly	Asn	Gly
	65					70					75				80	
	Thr	Ser	Asn	Lys	Gly	Met	Ala	Pro	Gln	Ala	Asn	Leu	Val	Phe	Gln	Ser
45				85					90						95	
	Val	Met	Asp	Ser	Asn	Gly	Gly	Leu	Gly	Gly	Leu	Pro	Ser	Asn	Val	Ser
			100						105					110		
50	Thr	Leu	Phe	Ser	Gln	Ala	Tyr	Ser	Ala	Gly	Ala	Arg	Ile	His	Thr	Asn
		115						120					125			
	Ser	Trp	Gly	Ala	Pro	Val	Asn	Gly	Ala	Tyr	Thr	Thr	Asp	Ser	Arg	Asn
55		130					135					140				

EP 1 209 233 A2

	Val	Asp	Asp	Tyr	Val	Arg	Lys	Asn	Asp	Met	Ala	Val	Leu	Phe	Ala	Ala	
	145					150					155					160	
5	Gly	Asn	Glu	Gly	Pro	Asn	Gly	Gly	Thr	Ile	Ser	Ala	Pro	Gly	Thr	Ala	
					165					170					175		
	Lys	Asn	Ala	Ile	Thr	Val	Gly	Ala	Thr	Glu	Asn	Leu	Arg	Pro	Ser	Phe	
10				180					185					190			
	Gly	Ser	Tyr	Ala	Asp	Asn	Ile	Asn	His	Val	Ala	Gln	Phe	Ser	Ser	Arg	
			195					200					205				
	Gly	Pro	Thr	Lys	Asp	Gly	Arg	Ile	Lys	Pro	Asp	Val	Met	Ala	Pro	Gly	
15		210					215					220					
	Thr	Phe	Ile	Leu	Ser	Ala	Arg	Ser	Ser	Leu	Ala	Pro	Asp	Ser	Ser	Phe	
	225					230					235					240	
20	Trp	Ala	Asn	His	Asp	Ser	Lys	Tyr	Ala	Tyr	Met	Gly	Gly	Thr	Ser	Met	
					245					250					255		
	Ala	Thr	Pro	Ile	Val	Ala	Gly	Asn	Val	Ala	Gln	Leu	Arg	Glu	His	Phe	
25				260					265					270			
	Ile	Lys	Asn	Arg	Gly	Ile	Thr	Pro	Lys	Pro	Ser	Leu	Leu	Lys	Ala	Ala	
			275					280					285				
	Leu	Ile	Ala	Gly	Ala	Thr	Asp	Ile	Gly	Leu	Gly	Tyr	Pro	Ser	Gly	Asn	
30		290					295					300					
	Gln	Gly	Trp	Gly	Arg	Val	Thr	Leu	Asp	Lys	Ser	Leu	Asn	Val	Ala	Phe	
	305					310					315					320	
	Val	Asn	Glu	Thr	Ser	Ser	Leu	Ser	Thr	Asn	Gln	Lys	Ala	Thr	Tyr	Ser	
35					325					330					335		
	Phe	Thr	Ala	Gln	Ser	Gly	Lys	Pro	Leu	Lys	Ile	Ser	Leu	Val	Trp	Ser	
				340					345					350			
40	Asp	Ala	Pro	Ala	Ser	Thr	Ser	Ala	Ser	Val	Thr	Leu	Val	Asn	Asp	Leu	
			355					360					365				
	Asp	Leu	Val	Ile	Thr	Ala	Pro	Asn	Gly	Thr	Lys	Tyr	Val	Gly	Asn	Asp	
45		370					375					380					
	Phe	Thr	Ala	Pro	Tyr	Asp	Asn	Asn	Trp	Asp	Gly	Arg	Asn	Asn	Val	Glu	
	385					390					395					400	
	Asn	Val	Phe	Ile	Asn	Ala	Pro	Gln	Ser	Gly	Thr	Tyr	Thr	Val	Glu	Val	
50				405						410					415		
	Gln	Ala	Tyr	Asn	Val	Pro	Gln	Gly	Pro	Gln	Ala	Phe	Ser	Leu	Ala	Ile	
				420				425						430			
55	Val	Asn															

EP 1 209 233 A2

<210> 7  
 <211> 433  
 <212> PRT  
 <213> Bacillus sp.  
  
 <400> 7  
 Asn Asp Val Ala Arg Gly Ile Val Lys Ala Asp Val Ala Gln Asn Asn  
 1 5 10 15  
 Phe Gly Leu Tyr Gly Gln Gly Gln Ile Val Ala Val Ala Asp Thr Gly  
 20 25 30  
 Leu Asp Thr Gly Arg Asn Asp Ser Ser Met His Glu Ala Phe Arg Gly  
 15 35 40 45  
 Lys Ile Thr Ala Leu Tyr Ala Leu Gly Arg Thr Asn Asn Ala Asn Asp  
 50 55 60  
 Pro Asn Gly His Gly Thr His Val Ala Gly Ser Val Leu Gly Asn Ala  
 20 65 70 75 80  
 Thr Asn Lys Gly Met Ala Pro Gln Ala Asn Leu Val Phe Gln Ser Ile  
 25 85 90 95  
 Met Asp Ser Gly Gly Gly Leu Gly Gly Leu Pro Ala Asn Leu Gln Thr  
 100 105 110  
 Leu Phe Ser Gln Ala Tyr Ser Ala Gly Ala Arg Ile His Thr Asn Ser  
 30 115 120 125  
 Trp Gly Ala Pro Val Asn Gly Ala Tyr Thr Thr Asp Ser Arg Asn Val  
 130 135 140  
 Asp Asp Tyr Val Arg Lys Asn Asp Met Thr Ile Leu Phe Ala Ala Gly  
 35 145 150 155 160  
 Asn Glu Gly Pro Gly Ser Gly Thr Ile Ser Ala Pro Gly Thr Ala Lys  
 165 170 175  
 Asn Ala Ile Thr Val Gly Ala Thr Glu Asn Leu Arg Pro Ser Phe Gly  
 40 180 185 190  
 Ser Tyr Ala Asp Asn Ile Asn His Val Ala Gln Phe Ser Ser Arg Gly  
 45 195 200 205  
 Pro Thr Arg Asp Gly Arg Ile Lys Pro Asp Val Met Ala Pro Gly Thr  
 210 215 220  
 Tyr Ile Leu Ser Ala Arg Ser Ser Leu Ala Pro Asp Ser Ser Phe Trp  
 50 225 230 235 240  
 Ala Asn His Asp Ser Lys Tyr Ala Tyr Met Gly Gly Thr Ser Met Ala  
 245 250 255  
 55

# EP 1 209 233 A2

Thr Pro Ile Val Ala Gly Asn Val Ala Gln Leu Arg Glu His Phe Val  
 260 265 270  
 5 Lys Asn Arg Gly Val Thr Pro Lys Pro Ser Leu Leu Lys Ala Ala Leu  
 275 280 285  
 Ile Ala Gly Ala Ala Asp Val Gly Leu Gly Phe Pro Asn Gly Asn Gln  
 290 295 300  
 10 Gly Trp Gly Arg Val Thr Leu Asp Lys Ser Leu Asn Val Ala Phe Val  
 305 310 315 320  
 Asn Glu Thr Ser Pro Leu Ser Thr Ser Gln Lys Ala Thr Tyr Ser Phe  
 15 325 330 335  
 Thr Ala Gln Ala Gly Lys Pro Leu Lys Ile Ser Leu Val Trp Ser Asp  
 340 345 350  
 20 Ala Pro Gly Ser Thr Thr Ala Ser Leu Thr Leu Val Asn Asp Leu Asp  
 355 360 365  
 Leu Val Ile Thr Ala Pro Asn Gly Thr Lys Tyr Val Gly Asn Asp Phe  
 370 375 380  
 25 Thr Ala Pro Tyr Asp Asn Asn Trp Asp Gly Arg Asn Asn Val Glu Asn  
 385 390 395 400  
 Val Phe Ile Asn Ala Pro Gln Ser Gly Thr Tyr Thr Val Glu Val Gln  
 30 405 410 415  
 Ala Tyr Asn Val Pro Val Ser Pro Gln Thr Phe Ser Leu Ala Ile Val  
 420 425 430  
 35 His

## Claims

- 40
1. An alkaline protease wherein an amino acid residue at (a) position 84, (b) position 104, (c) position 256 or (d) position 369 of SEQ ID NO:1 or at a position corresponding thereto has been deleted or selected from:
 

45

    - at position (a): an arginine residue,
    - at position (b): a proline residue,
    - at position (c): an alanine, serine, glutamine, valine, leucine, asparagine, glutamic acid or aspartic acid residue,
    - and
    - at position (d): an aspartic acid residue.
  2. An alkaline protease having an amino acid sequence represented by SEQ ID NO:1 or having an amino acid sequence showing at least 60% homology therewith, wherein an amino acid residue at (a) position 84, (b) position 104, (c) position 256 or (d) position 369 of SEQ ID NO:1 or at a position corresponding thereto has been deleted or selected from:
 

55

    - at position (a): an arginine residue,
    - at position (b): a proline residue,
    - at position (c): an alanine, serine, glutamine, valine, leucine, asparagine, glutamic acid or aspartic acid residue,
    - and

at position (d): an aspartic acid residue.

3. An alkaline protease wherein an amino acid residue at (e) position 66 or 264, (f) position 57, each of 101 to 106, 136, 193 or 342, (g) position 46 or 205, (h) position 54, 119, 138, 148 or 195, (i) position 247, (j) position 124, (k) position 107 or (l) position 257 of SEQ ID NO:1, or at a position corresponding thereto has been deleted or selected from::

at position (e): a glutamine, aspartic acid, serine, glutamic acid, alanine, threonine, leucine, methionine, cysteine, valine, glycine or isoleucine residue

at position (f): a lysine, serine, glutamine, phenylalanine, valine, arginine, tyrosine, leucine, isoleucine, threonine, methionine, cysteine, tryptophan, aspartic acid, glutamic acid, histidine, proline or alanine residue,

at position (g): a tyrosine, tryptophan, alanine, asparagine, glutamic acid, threonine, valine, leucine, isoleucine, histidine, serine, lysine, glutamine, methionine or cysteine residue,

at position (h): a tryptophan, phenylalanine, alanine, asparagine, glutamic acid, threonine, valine, histidine, serine, lysine, glutamine, methionine, glycine, aspartic acid, proline, arginine or cysteine residue,

at position (i): a tryptophan, phenylalanine, alanine, asparagine, glutamic acid, threonine, valine, leucine, isoleucine, histidine, serine, glutamine, methionine or cysteine residue,

at position (j): an alanine or lysine residue,

at position (k): a lysine, arginine, alanine or serine residue, and

at position (l): a valine or isoleucine residue.

4. An alkaline protease having an amino acid sequence represented by SEQ ID NO:1 or having an amino acid sequence showing at least 60% homology therewith, wherein an amino acid residue at (e) position 66 or 264, (f) position 57, each of 101 to 106, 136, 193 or 342, (g) position 46 or 205, (h) position 54, 119, 138, 148 or 195, (i) position 247, (j) position 124, (k) position 107 or (l) position 257 has been deleted or selected from:

at position (e): a glutamine, aspartic acid, serine, glutamic acid, alanine, threonine, leucine, methionine, cysteine, valine, glycine or isoleucine residue

at position (f): a lysine, serine, glutamine, phenylalanine, valine, arginine, tyrosine, leucine, isoleucine, threonine, methionine, cysteine, tryptophan, aspartic acid, glutamic acid, histidine, proline or alanine residue,

at position (g): a tyrosine, tryptophan, alanine, asparagine, glutamic acid, threonine, valine, leucine, isoleucine, histidine, serine, lysine, glutamine, methionine or cysteine residue,

at position (h): a tryptophan, phenylalanine, alanine, asparagine, glutamic acid, threonine, valine, histidine, serine, lysine, glutamine, methionine, glycine, aspartic acid, proline, arginine or cysteine residue,

at position (i): a tryptophan, phenylalanine, alanine, asparagine, glutamic acid, threonine, valine, leucine, isoleucine, histidine, serine, glutamine, methionine or cysteine residue,

at position (j): an alanine or lysine residue,

at position (k): a lysine, arginine, alanine or serine residue, and

at position (l): a valine or isoleucine residue.

5. An alkaline protease according to claim 2 or 4, wherein the amino acid sequence represented by SEQ ID NO:1 or amino acid sequence showing at least 60% homology therewith is an amino acid sequence selected from SEQ ID NOS: 2 to 7.

6. A gene encoding an alkaline protease as claimed in any one of claims 1 to 5.


7. A recombinant vector comprising a gene as claimed in claim 6.

8. A transformant comprising a recombinant vector as claimed in claim 7.

9. A transformant according to claim 8, wherein a microorganism is used as a host.


10. A detergent composition comprising an alkaline protease as claimed in any one of claims 1 to 5.

FIG. 1



BEST AVAILABLE COPY

FIG. 2



BEST AVAILABLE COPY

FIG. 3

